



Contribution ID : 190

Type : **Oral**

Understanding the Dynamics of the Oxic-Anoxic Interface in the Black Sea

Wednesday, 5 September 2018 17:15 (15)

The Black Sea, the largest semienclosed anoxic basin on Earth, can be considered as an excellent natural laboratory for oxic and anoxic biogeochemical processes. The suboxic zone, a thin interface between oxic and anoxic waters, still remains poorly understood because it has been undersampled. This has led to alternative concepts regarding the underlying processes that create it. Existing hypotheses suggest that the interface originates either by isopycnal intrusions that introduce oxygen or the dynamics of manganese redox cycling that are associated with the sinking of particles or chemosynthetic bacteria. Here we reexamine these concepts using high-resolution oxygen, sulfide, nitrate, and particle concentration profiles obtained with sensors deployed on profiling floats. Our results show an extremely stable structure in density space over the entire basin with the exception of areas near the Bosphorus plume and in the southern areas dominated by coastal anticyclones. The absence of large-scale horizontal intrusive signatures in the open-sea supports a hypothesis prioritizing the role of biogeochemical processes.

Position

Professor

Affiliation

1Institute of Coastal Research, Helmholtz-Zentrum Geesthacht, Geesthacht, Germany

Email Address

emil.stanev@hzg.de

Are you a SFB 754 / Future Ocean member?

No

Primary author(s) : STANEV, Emil Vassilev (1Institute of Coastal Research, Helmholtz-Zentrum Geesthacht)

Presenter(s) : STANEV, Emil Vassilev (1Institute of Coastal Research, Helmholtz-Zentrum Geesthacht)

Session Classification : 03 Ventilation and Oxygen Supply

Track Classification : 03 Ventilation and Oxygen Supply